

Possibly New Charmed Baryon States from $\bar{B}^0 \rightarrow p\bar{p}D^{(*)0}$ Decays

Hai-Yang Cheng,¹ C.Q. Geng,^{2,3} and Y.K. Hsiao³

¹*Institute of Physics, Academia Sinica, Taipei, Taiwan 115, R.O.C.*

²*Department of Physics, National Tsing Hua University, Hsinchu, Taiwan 300, R.O.C.*

³*Physics Division, National Center for Theoretical Sciences, Hsinchu, Taiwan 300, R.O.C.*

(Dated: March 3, 2013)

Abstract

Based on the perturbative QCD counting rules, we fit the invariant mass spectra of $D^{(*)0}p$ in $\bar{B}^0 \rightarrow p\bar{p}D^{(*)0}$ decays measured by BABAR. We find that some peaks in the spectra arise from the charmed baryon states denoted by $\mathbf{B}_c$ through the 2-step processes of $\bar{B}^0 \rightarrow \mathbf{B}_c^+(\rightarrow D^{(*)0}p)\bar{p}$. In addition to the established states $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$, two of them, namely, $\mathbf{B}_c(2820)^+$ and $\mathbf{B}_c(3280)^+$, may be already revealed by experiments, while the rest, identified as $\mathbf{B}_c(3000)^+$, $\mathbf{B}_c(3100)^+$, $\mathbf{B}_c(3120)^+$, $\mathbf{B}_c(3200)^+$, and $\mathbf{B}_c(3320)^+$, can be viewed as an indication of new heavy charmed baryons.

PACS numbers: 14.20.Lq, 13.85.Ni, 13.25.Hw, 12.38.Bx

Introduction— The charmed baryon spectroscopy is expected to be an ideal place to study the dynamics of the light quarks in the environment of a heavy quark. Apart from a charm quark, the charmed baryon contains two light quarks as a diquark, which can be either an antitriplet or a sextet in the quark model. Due to various combinations of individual spins and orbital angular momenta between two light quarks as well as the diquark and charm quark, charmed baryons exhibit a rich spectrum of states [1–4]. However, so far only a few of charmed baryons have been experimentally identified [5, 6]. The B factories as well as the LHCb are able to play an important role in studying charmed baryons. For example, the charmed baryon states of $\Lambda_c(2665)^+$, $\Lambda_c(2880)^+$, $\Lambda_c(2940)^+$, $\Sigma_c(2800)$, $\Xi_c(2980)$, and $\Xi_c(3080)$, have been observed by CLEO [7], BELLE [8], and BABAR [9]. In particular, the decays of $\Lambda_c(2880)^+ \rightarrow \Sigma_c(2520)^{0,++}\pi^{+,-}$ have been analyzed to assign the quantum numbers of $\Lambda_c(2880)^+$ to be $J^P = \frac{5}{2}^+$ [10].

Recently, the BABAR collaboration has reported that both $m_{p\bar{p}}$ and $m_{D^{(*)}0p}$ invariant mass distributions in $\bar{B}^0 \rightarrow p\bar{p}D^{(*)0}$ decays exhibit low-mass enhancement in the spectra [11]. In the $m_{p\bar{p}}$ spectra, the low-mass peaks near threshold can be recognized as the threshold effect [12]. On the one hand, the exclusion for the ranges of $m_{D^0p}^2 < 9 \text{ GeV}^2$ and $m_{D^{*0}p}^2 < 10.5 \text{ GeV}^2$ improves the knowledge of the threshold effect, a generic feature in three-body baryonic $\bar{B} \rightarrow \mathbf{B}\bar{\mathbf{B}}'M$ decays. On the other hand, the low-mass peaks in the $m_{D^{(*)}p}$ spectra with $m_{p\bar{p}}^2 > 5 \text{ GeV}^2$ seem puzzling as the data points disagree with what expected from the uniform phase-space model. In this letter, we find that, unlike the peaks around the threshold area in the $m_{p\bar{p}}$ spectra, the low-mass peaks in the $m_{D^{(*)}0p}$ spectra can be understood based on the pQCD effect in the creation of the dibaryon [13, 14]. We will demonstrate that the charmed-baryon resonances decaying into $D^{(*)0}p$ fit well to these peaking data points in $\bar{B}^0 \rightarrow p\bar{p}D^{(*)0}$ decays. It turns out that some of the resonant charmed baryon states are not experimentally observed yet, but theoretically studied in the literature [1–4].

Formalism— To explain the low-mass enhancement in the $m_{D^{(*)}0p}$ invariant mass distributions in $\bar{B}^0 \rightarrow p\bar{p}D^{(*)0}$, the relevant simplified reduced amplitudes are given by [14]

$$\begin{aligned}\mathcal{A}(\bar{B}^0 \rightarrow p\bar{p}D^0) &\simeq -\frac{G_F}{\sqrt{2}}V_{cb}V_{ud}^*a_2f_Dm_D^2\bar{u}_p(f_3 + g_3\gamma_5)v_{\bar{p}}, \\ \mathcal{A}(\bar{B}^0 \rightarrow p\bar{p}D^{*0}) &\simeq i\frac{G_F}{\sqrt{2}}V_{cb}V_{ud}^*a_2m_{D^*}f_{D^*}\sigma_{\mu\nu}\varepsilon^{\mu*}p^\nu\bar{u}_p(f_2 + g_2\gamma_5)v_{\bar{p}},\end{aligned}\quad (1)$$

with the Fermi constant G_F , the CKM matrix elements $V_{q_iq_j}$, and $p = p_{\bar{B}^0} - (p_p + p_{\bar{p}})$. Here, a_2 is a parameter that accommodates the nonfactorizable effects, the decay constants $f_{D^{(*)}}$

and the polarization 4-vector $\varepsilon^{\mu*}$ come from the matrix elements of $\langle D|\bar{c}\gamma^\mu\gamma_5 u|0\rangle = -if_D p^\mu$ and $\langle D^*|\bar{c}\gamma^\mu u|0\rangle = m_{D^*}f_{D^*}\varepsilon^{\mu*}$, and the form factors f_i and g_i with $i=2$ and 3 can be traced back to the parameterization for the matrix elements of the $\bar{B} \rightarrow \mathbf{B}\bar{\mathbf{B}}'$ transitions [14, 15], of which the momentum dependences are given by

$$f_i = \frac{D_{f_i}}{t^3}, \quad g_i = \frac{D_{g_i}}{t^3}, \quad (2)$$

with $t \equiv m_{p\bar{p}}^2$ and the constants D_{f_i} and D_{g_i} . According to pQCD counting rules [16, 17], the momentum dependence of the form factors $f_{2,3}$ and $g_{2,3}$ is proportional to $1/t^3$ as three hard gluons are needed to produce $\mathbf{B}\bar{\mathbf{B}}'$. Since the $1/t^3$ function peaks at the low mass and flattens out at the large energy region, the threshold effect in the $m_{p\bar{p}}$ spectra [14] can be easily accommodated. Nonetheless, as we shall see, the amplitudes in Eq. (1) have no other short-distance contributions to explain the low-mass peaking points in the $m_{D^{(*)0}p}$ spectra. Inspired by the observations of $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+ \rightarrow D^0 p$ [9], we shall examine the $\bar{B}^0 \rightarrow p\bar{p}D^{(*)0}$ decays via the resonant $\bar{B}^0 \rightarrow \mathbf{B}_c(\rightarrow D^{(*)0}p)\bar{p}$ channels to explain the $m_{D^{(*)0}p}$ invariant mass spectra, where $\mathbf{B}_c$ represents the possible charmed baryon states. Accordingly, the resonant amplitudes read

$$\mathcal{A}_{\mathcal{R}}(\bar{B}^0 \rightarrow p\bar{p}D^{(*)0}) = \frac{G_F}{\sqrt{2}} V_{cb} V_{ud}^* \bar{u}_p \mathcal{R} (a + b\gamma_5) v_{\bar{p}}, \quad (3)$$

where the factor $\mathcal{R}$ is defined by

$$\mathcal{R} = \frac{\not{q} + m}{q^2 - m^2 + im\Gamma}, \quad (4)$$

with $q = p_p + p_{D^{(*)0}}$ and m (Γ) the mass (width) of the resonance. Note that the parameters a and b in Eq. (3) may contain the combinations of 4-vector indices. Besides, even if $\mathbf{B}_c$ carries a higher spin, such as $\Lambda_c(2880)^+$ with spin-5/2, the spin structure in the propagator can still be summed over and factored into a and b . Since the couplings of $\mathbf{B}_c \rightarrow D^{(*)0}p$ and the parity of $\mathbf{B}_c$ may not be well determined, there exist some other possibilities for the amplitudes in Eq. (3), such as $(a - b\gamma_5)$ and $(b \pm a\gamma_5)$. By taking $|a| = |b|$, we are able to circumvent this complexity. Hence, we can obtain the amplitude squared $|\bar{\mathcal{A}}|^2$ with the bar denoting the summation over p and $\bar{p}$ spins. Using the general partial decay rate formula for the 3-body decay given by [5]

$$d\Gamma = \frac{1}{(2\pi)^3} \frac{|\bar{\mathcal{A}}|^2}{32M_{\bar{B}^0}^3} dm_{D^{(*)0}p}^2 dm_{p\bar{p}}^2, \quad (5)$$




FIG. 1. Invariant mass spectra as functions of the invariant masses $m_{D^0 p}$ and $m_{D^{*0} p}$ in $\bar{B}^0 \rightarrow p\bar{p}D^0$ and $\bar{B}^0 \rightarrow p\bar{p}D^{*0}$ decays, respectively, where the solid lines include both the resonant and non-resonant contributions, while the dashed lines correspond to the non-resonant contributions only.

we can integrate over $m_{p\bar{p}}^2$ to study the partial rates as functions of $m_{D^{(*)0}p}$ in $\bar{B}^0 \rightarrow p\bar{p}D^{(*)0}$.

Numerical Analysis— To illustrate our results, in our numerical analysis we take the CKM matrix elements $V_{cb} = A\lambda^2$ and $V_{ud} = 1 - \lambda^2/2$ with $A = 0.808$ and $\lambda = 0.2253$ from the Particle Data Group [5]. The decay constants of $D^{(*)}$ are given as $(f_D, f_{D^*}) = (0.22, 0.23)$ GeV [5, 18]. The parameter $a_2 = 0.33 \pm 0.04$ is extracted from the total branching fractions of $\bar{B}^0 \rightarrow p\bar{p}D^{(*)0}$. The constants D_{f_i} and D_{g_i} in Eq. (2) have been fitted in Ref. [14]. Explicitly, we adopt their values to be

$$D_{g_2} = -D_{f_2} = (62.4 \pm 8.9) \text{ GeV}^4, \quad D_{g_3} = -D_{f_3} = (280.0 \pm 44.0) \text{ GeV}^4. \quad (6)$$

Because the low-mass enhancements in the $m_{D^{(*)0}p}$ invariant mass spectra are observed for $m_{p\bar{p}}^2 > 5 \text{ GeV}^2$, we integrate $m_{p\bar{p}}^2$ in Eq. (5) over the range of $5 \text{ GeV}^2 < m_{p\bar{p}}^2 < (m_{p\bar{p}}^2)_{max}$ with $(m_{p\bar{p}}^2)_{max}$ as a function of $m_{D^{(*)0}p}$ defined in Ref. [5]. As seen in Fig. 1, we find that, in spite of failing to peak at the low mass, the dashed lines due to the nonresonant amplitudes in Eq. (1) agree well with the non-peaking data points. On the contrary, these non-peaking data points are not in accordance with the shaded regions presented in Ref. [10] with the normalized data in the uniform phase-space model. Since the amplitudes in Eq. (1) originating from the short-distance contributions can reliably explain the non-peaking points, it is natural to describe the low-mass peaking points in terms of the resonant $\mathbf{B}_c$ decaying into $D^{(*)0}p$, and fit these points by adding sufficient amplitudes in the form of $\mathcal{A}_R$ in Eq. (3). Again, as shown in Fig. 1, to smoothly link all the peaking data points with the solid lines, we

need six and three charmed baryons as resonant states in the $\bar{B}^0 \rightarrow p\bar{p}D^0$ and $\bar{B}^0 \rightarrow p\bar{p}D^{*0}$ decays, respectively. Being named from their fit masses, these charmed baryons from left to right in the m_{D^0p} and $m_{D^{*0}p}$ invariant mass spectra are orderly listed in Table I.

In the m_{D^0p} spectrum, the second $\mathbf{B}_c$ with the fit mass and decay width $(m, \Gamma) = (2880, 6)$ MeV and the third $\mathbf{B}_c$ with $(m, \Gamma) = (2940, 24)$ MeV have been identified as the experimentally confirmed charmed baryons $\Lambda_c(2880)^+$ with $(m, \Gamma) = (2881.53 \pm 0.35, 5.8 \pm 1.1)$ MeV and $\Lambda_c(2940)^+$ with $(m, \Gamma) = (2939.3^{+1.4}_{-1.5}, 17^{+8}_{-6})$ MeV, respectively. In Ref. [9] there is a suspected resonance around 2840 MeV, which may relate to $\mathbf{B}_c(2820)^+$. Moreover, according to the BABAR's preliminary result in the study of $B^- \rightarrow \Sigma_c^{++}\bar{p}\pi^-\pi^-$ [19], the unexplained resonance with $(m, \Gamma) = (3245 \pm 20, 108 \pm 6)$ MeV in the spectrum of $\Sigma_c^{++}\pi^-\pi^-$ could be the neutral component of the excited baryon states $(\Sigma_c^{++}, \Sigma_c^+, \Sigma_c^0)$, in which the charge one may be in accordance with $\mathbf{B}_c(3280)^+$. Note that this type of the states has been predicted in Ref. [3]. It is interesting to point out that most of the charmed baryons listed in Table I can find their places in the theoretically predicted charmed baryon spectrum [1–4]. Note that $\mathbf{B}_c(2820)^+$ cannot be a $\Sigma_c(2800)^+$ as it is below the threshold of D^0p . By comparing the fit masses in Table I with the predicted ones in Ref. [1], the state $\mathbf{B}_c(2820)^+$ with $(m, \Gamma) = (2820, 5)$ MeV can be a Σ_c with $m = 2817$ MeV and $J^P = 1/2^-$, whereas the other Σ_c with $m = 2815$ MeV and $J^P = 3/2^-$ seems to be consistent with $\Sigma_c(2800)^+$ [2]. Since there are many theoretical predicted $\mathbf{B}_c$ states of order 3000 MeV [3], it may not be easy to identify $\mathbf{B}_c(3000)^+$ in the $m_{D^{*0}p}$ spectrum. As shown in Table I, because of the tiny mass difference and the same decay width, $\mathbf{B}_c(3100)^+$ and $\mathbf{B}_c(3120)^+$ are likely to be the same

TABLE I. Possible charmed baryons ($\mathbf{B}_c$) as the resonant states in the decays of $\bar{B}^0 \rightarrow p\bar{p}D^0$ (left panel) and $\bar{B}^0 \rightarrow p\bar{p}D^{*0}$ (right panel) with the units of $|a|$ and $|b|$ in GeV^4 .

$\mathbf{B}_c$ State	$ a = b $	(m, Γ) MeV	$\mathbf{B}_c$ State	$ a = b $	(m, Γ) MeV
$(2820)^+$	0.033	(2820, 5)	$(3000)^+$	0.014	(3000, 10)
$\Lambda_c(2880)^+$	0.009	(2880, 6)	$(3100)^+$	0.005	(3100, 10)
$\Lambda_c(2940)^+$	0.038	(2940, 24)	$(3280)^+$	0.020	(3280, 90)
$(3120)^+$	0.003	(3120, 10)			
$(3200)^+$	0.040	(3200, 65)			
$(3320)^+$	0.005	(3320, 20)			

particle, and the identification can be one of the excited Λ_c baryons with the (1F) or (3S) orbital excitation of the c-(ud) bound state [3]. On the other hand, the state $\mathbf{B}_c(3200)^+$ can be identified as one or the combination of the excited Σ_c baryons with masses of 3176, 3180, and 3186 MeV, coming from the orbital (2P) excitations in the c-(ud) bound state [3]. With the heaviest fit masses in the $m_{D^0 p}$ and $m_{D^{*0} p}$ spectra, the charmed baryon states $\mathbf{B}_c(3280)^+$ and $\mathbf{B}_c(3320)^+$, respectively, can be related to one or the combination of Σ_c with $m = 3262$ MeV and $J^P = 3/2^+$ and Σ_c with $m = 3268$ MeV and $J^P = 5/2^+$ from the orbital (2D) excitations of the c-(ud) bound state [3].

As for the resonant charmed baryon states found in $\bar{B}^0 \rightarrow p\bar{p}D^{(*)0}$ decays, one can perform experimental searches. In analog to the first observation of $\Lambda_c(2940)^+$ in the process of $e^+e^- \rightarrow \Lambda_c(2940)^+(\rightarrow D^0 p)X$ by BABAR [9], the $\mathbf{B}_c$ states can be found by scanning the $D^{(*)0}p$ spectra in $\mathbf{B}_c \rightarrow D^{(*)0}p$ from the e^+e^- , $p\bar{p}$, and pp colliders at the B factories, Tevatron, and LHC, respectively. We also expect that some of the resonant B_c states should be observed in the B decays through $\bar{B}^0 \rightarrow \mathbf{B}_c^+(\rightarrow \Sigma_c(2455)^{0,++}\pi^\pm)\bar{p}$, $\bar{B}^0 \rightarrow \mathbf{B}_c^+(\rightarrow \Sigma_c(2520)^{0,++}\pi^\pm)\bar{p}$, $B^- \rightarrow \mathbf{B}_c^+(\rightarrow \Sigma_c(2455)^{0,++}\pi^\pm)\bar{p}\pi^-$, and $B^- \rightarrow \mathbf{B}_c^+(\rightarrow \Sigma_c(2520)^{0,++}\pi^\pm)\bar{p}\pi^-$. It is important to note that as the resonant $\bar{B}^0 \rightarrow p\bar{p}D^{(*)0}$ decays imply the existence of the two-body $\bar{B}^0 \rightarrow \mathbf{B}_c^+\bar{p}$ decays, e.g. $\bar{B}^0 \rightarrow \Lambda_c(2940)^+\bar{p}$, the angular analysis in the $\bar{B}^0 \rightarrow \mathbf{B}_c^+\bar{p}$ decays will be very useful to assign the right quantum numbers to these $\mathbf{B}_c$ states.

Conclusions— We have analyzed the low-mass peaks in the $m_{D^{(*)0}p}$ spectra with $m_{p\bar{p}}^2 > 5 \text{ GeV}^2$ observed in $\bar{B}^0 \rightarrow p\bar{p}D^{(*)0}$ decays. We have found that the peaking points can be recognized as the charmed baryon resonances in $\bar{B}^0 \rightarrow \mathbf{B}_c(\rightarrow D^{(*)0}p)\bar{p}$ channels. Explicitly, we have introduced six charmed baryon states in the $m_{D^0 p}$ spectrum and three in $m_{D^{*0} p}$, in which two of them are readily identified as $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$. For other charmed baryon states, the current experimental data may have hinted at the existence of $\mathbf{B}_c(2820)^+$ and $\mathbf{B}_c(3280)^+$, while $\mathbf{B}_c(3000)^+$, $\mathbf{B}_c(3100)^+$, $\mathbf{B}_c(3120)^+$, $\mathbf{B}_c(3200)^+$, $\mathbf{B}_c(3320)^+$ can be referred to the theoretically predicted states in the literature, which could be discovered by various collider searches soon. Moreover, since the resonant $\bar{B}^0 \rightarrow \mathbf{B}_c(\rightarrow D^{(*)0}p)\bar{p}$ decays are the consequences of the two-body $\bar{B}^0 \rightarrow \mathbf{B}_c^+\bar{p}$ decays, the studies of the angular distributions in the two-body modes will help us to determine these new charmed baryon states.

Acknowledgments— We would like to thank Dr. Yubing Dong for useful discussions. The work was supported in part by National Center of Theoretical Science, and National Science

- [1] S. Migura, D. Merten, B. Metsch and H. R. Petry, Eur. Phys. J. A **28**, 41 (2006).
- [2] H.Y. Cheng, in *Proceedings of International Workshop on Charm Physics (Charm 2007)*, Ithaca, New York, 5-8 Aug 2007; H.Y. Cheng and C. K. Chua, Phys. Rev. D **75**, 014006 (2007).
- [3] D. Ebert, R. N. Faustov and V. O. Galkin, Phys. Lett. B **659**, 612 (2008); Phys. Rev. D **84**, 014025 (2011).
- [4] C. Chen, X. L. Chen, X. Liu, W. Z. Deng and S. L. Zhu, Phys. Rev. D **75**, 094017 (2007); A. Valcarce, H. Garcilazo and J. Vijande, Eur. Phys. J. A **37**, 217 (2008); D. Gamermann, C. Garcia-Recio, J. Nieves, L. L. Salcedo and L. Tolos, Phys. Rev. D **81**, 094016 (2010); O. Romanets, L. Tolos, C. Garcia-Recio, J. Nieves, L. L. Salcedo and R. G. E. Timmermans, arXiv:1202.2239 [hep-ph].
- [5] K. Nakamura *et al.* [Particle Data Group], J. Phys. G **37**, 075021 (2010).
- [6] E. Klempt and J. M. Richard, Rev. Mod. Phys. **82**, 1095 (2010).
- [7] M. Artuso *et al.* [CLEO Collaboration], Phys. Rev. Lett. **86**, 4479 (2001).
- [8] R. Mizuk *et al.* [BELLE Collaboration], Phys. Rev. Lett. **94**, 122002 (2005); R. Chistov *et al.* [BELLE Collaboration], Phys. Rev. Lett. **97**, 162001 (2006).
- [9] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **98**, 012001 (2007).
- [10] K. Abe *et al.* [BELLE Collaboration], Phys. Rev. Lett. **98**, 262001 (2007).
- [11] P. del Amo Sanchez *et al.* [The BABAR Collaboration], arXiv:1111.4387 [hep-ex].
- [12] W.S. Hou and A. Soni, Phys. Rev. Lett. **86**, 4247 (2001).
- [13] Y.K. Hsiao, Int. J. Mod. Phys. A **24**, 3638 (2009); C.Q. Geng and Y.K. Hsiao, Phys. Lett. B **704**, 495 (2011); Phys. Rev. D **85**, 017501 (2012).
- [14] C.H. Chen, H.Y. Cheng, C.Q. Geng and Y.K. Hsiao, Phys. Rev. D **78**, 054016 (2008).
- [15] C.Q. Geng and Y.K. Hsiao, Phys. Rev. D **74**, 094023 (2006); C.Q. Geng, Y.K. Hsiao and J.N. Ng, Phys. Rev. Lett. **98**, 011801 (2007); Phys. Rev. D **75**, 094013 (2007).
- [16] G.P. Lepage and S.J. Brodsky, Phys. Rev. Lett. **43**, 545(1979) [Erratum-ibid. **43**, 1625 (1979)]; Phys. Rev. D **22**, 2157 (1980); S.J. Brodsky, G.P. Lepage and S. A. A. Zaidi, Phys. Rev. D

23, 1152 (1981).

[17] C.K. Chua, W.S. Hou and S. Y. Tsai, Phys. Rev. D **66**, 054004 (2002).

[18] H.M. Choi, Phys. Rev. D **75**, 073016 (2007).

[19] Talk presented by O. Grüberg for the BABAR Collaboration at the conference of the XLVIIth Rencontres de Moriond, La Thuile, March 10-17, 2012.